

Armor  **Tech**

ZOXY 2F

Use as a broad spectrum fungicide for control of listed diseases on labeled turf sites.

ACTIVE INGREDIENTS

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate*.....	22.9%
OTHER INGREDIENTS:	77.1%
TOTAL:	100.00%

Contains 2.08 pounds of azoxystrogin per gallon.

**IUPAC*

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No: 86064-6
EPA Est. No: 67545-AZ-001

NET CONTENTS: 1 QT

Manufactured for:
United Turf Alliance, LLC
8014 Cumming Highway, Suite 403-282
Canton, GA 30115



FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing should contact a physician.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates.

Azoxystrobin can be persistent for several months or longer.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and United Turf Alliance, LLC immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Failure to follow the use directions and precautions on this label may result in plant injury or poor disease control.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USES

Commercial turf farm use (Not for use in California).

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks

NON-AGRICULTURAL USES

Golf Courses (Not for use in California).

For use to control diseases on turf on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with this product is dry.

PRODUCT INFORMATION

ArmorTech ZOXY 2F is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. ArmorTech ZOXY 2F may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

RESTRICTIONS

Do not graze or feed clippings from treated turf areas to animals.

DO NOT spray ArmorTech ZOXY 2F where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply ArmorTech ZOXY 2F to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

PRECAUTIONS

ArmorTech ZOXY 2F is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

ArmorTech ZOXY 2F may demonstrate some phytotoxic effects when mixed with products that are formulated as ECs. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification is recommended.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of ArmorTech ZOXY 2F has been used. If resistant isolates to Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under certain conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

INTEGRATED PEST (DISEASE) MANAGEMENT

ArmorTech ZOXY 2F should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. This should include selection of varieties with disease tolerance, removal of plant debris in which inoculums overwinters, and proper timing and placement of irrigation. Consult your local agricultural authorities for additional IPM strategies established for your area. ArmorTech ZOXY 2F may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

GROUP 11 FUNGICIDE

ArmorTech ZOXY 2F (azoxystrobin) is a Group 11 fungicide. The mode of action for ArmorTech ZOXY 2F is the inhibition of the QoI (quinone outside) site within the electron transport system [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance develop cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include alternating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. United Turf Alliance, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the crop specific resistance management recommendations in the directions for use.

If no resistance recommendation on number of applications is specified in the directions for use, follow the recommendations in the table below.

If planned total number of fungicide applications per crop is:	1	2	3	4	5	6	7	8	9	10	11	12
Recommended Solo QoI fungicide sprays	1	1	2	2	2	2	2	3	3	3	3	4
Recommended QoI fungicide sprays in mixture (tank-mix formulated)	1	2	2	2	2	3	3	4	4	5	5	6

In situations requiring multiple sprays, develop season long spray programs for Group 11 (QoI) fungicides. In crops where two sequential Group 11 fungicide applications are made, they should be alternated with two or more applications of a fungicide that is not in Group 11. If more than 12 applications are made, observe the following guidelines:

- When using QoI fungicide as a solo product, the number of applications must be no more than $\frac{1}{3}$ (33%) of the total number of fungicide applications per season.
- For QoI mixes in programs in which tank mixes or pre mixes of QoI with mixing partners of a different mode of action are utilized, the number of QoI containing applications must be no more than $\frac{1}{2}$ (50%) of the total number of fungicide applications per season.
- In programs in which applications of QoI are made with both solo products and mixtures, the number of QoI containing applications must be no more than $\frac{1}{2}$ (50%) of the total number of fungicide applications per season.

If Group 11 fungicide is applied to the seed or soil, do not make another application with a Group 11 fungicide for at least 3 weeks.

SPRAY DRIFT MANAGEMENT

To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

ATTENTION

ArmorTech ZOXY 2F is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray ArmorTech ZOXY 2F where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply ArmorTech ZOXY 2F to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

MIXING AND APPLICATION METHODS

Spray Equipment

ArmorTech ZOXY 2F may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on the suction side of the pump should be 16-mesh or coarser.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.the total number of fungicide applications per season.

Pump

- Use a pump with capacity to:
 - (1) Maintain 35-40 psi at nozzles
 - (2) Provide sufficient agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

Mixing Instructions

- ArmorTech ZOXY 2F is a suspension concentrate (SC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

ArmorTech Zoxy 2F Alone (No Tank Mix)

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add ArmorTech ZOXY 2F to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after ArmorTech ZOXY 2F has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

ArmorTech Zoxy 2F + Tank Mixtures: ArmorTech ZOXY 2F is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of ArmorTech ZOXY 2F with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

ArmorTech ZOXY 2F has demonstrated some phytotoxic effects when mixed with products that are formulated as emulsifiable concentrates (EC). These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

Mixing in the Spray Tank

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and ArmorTech ZOXY 2F to the spray tank.
- Allow ArmorTech ZOXY 2F to completely disperse.
- Spray the mixture with the agitator running.

TURF

Not approved for use on Turf in California

ArmorTech ZOXY 2F is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), Fusarium patch, gray leaf spot, gray snow mold (Typhula blight), leafspot, melting out, necrotic ring spot, pink patch, pink snow mold, Pythium blight, Pythium root rot, red thread, Rhizoctonia large patch, southern blight, spring dead spot, summer patch, take-all patch, and Zoysia patch on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Integrated Pest (Disease) Management: Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Resistance Management: Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. ArmorTech ZOXY 2F should be applied in a tank mix or alternation program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not apply more than two sequential ArmorTech ZOXY 2F applications for Pythium spp. control. For all other diseases when Pythium spp. is not present, do not apply more than three sequential applications of ArmorTech ZOXY 2F.

Application Directions: ArmorTech ZOXY 2F should be applied prior to disease development. Mix ArmorTech ZOXY 2F with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 fl. oz. ArmorTech ZOXY 2F per 1 to 2 gallons of water. Do not apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1000 square feet/year). Apply by ground only.

Rate Ranges: Use the shortest specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

Dollar Spot: ArmorTech ZOXY 2F does not control dollar spot. ArmorTech ZOXY 2F is compatible in tank mixes with many other fungicides that control dollar spot. Always tank mix ArmorTech ZOXY 2F with another fungicide that controls dollar spot when this disease is present. Follow directions under TANK MIXES/COMPATIBILITY above.

DIRECTIONS FOR APPLICATION FOR TURF DISEASES

Targeted Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (days)	Remarks*
Anthracnose (<i>Colletotrichum graminicola</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Brown Patch (<i>Rhizoctonia solani</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Cool weather brown patch Yellow patch (<i>Rhizoctonia cerealis</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development.
Fusarium patch (<i>Microdochium nivale</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.38-0.77	14-28	Begin applications before disease is present and continue applications while conditions are favorable for disease development.
Gray snow mold Typhula blight (<i>Typhula incarnata</i> , <i>T. ishikariensis</i>)	1.35 0.77	Single application 14	Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure.
Leafspot (<i>Bipolaris sorokiniana</i>)	0.38-0.77	14-21	Apply when conditions are favorable for disease development.
Melting out (<i>Drechslera poae</i>)	0.38-0.77	14-21	Apply when conditions are favorable for disease development.
Necrotic ring spot (<i>Leptosphaeria korrae</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Pink patch (<i>Limonomyses roseipellis</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Pink snow mold (<i>Microdochium nivale</i>)	1.35 0.77	Single application 14	Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure.

Targeted Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (days)	Remarks*
Pythium blight Pythium root rot (<i>Pythium aphanidermatum</i> , <i>Pythium</i> spp.)	0.38-0.77	10-14	Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded as well as established turf.
Red thread (<i>Laetisaria fuciformis</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Rhizoctonia large patch (<i>Rhizoctonia solani</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development
Southern blight (<i>Sclerotium rolfsii</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Spring dead spot (<i>Leptosphaeria korrae</i>) or (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) or (<i>Ophiosphaerella herpotricha</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development.
Summer patch (<i>Magnaporthe poae</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Take-all patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.38-0.77	28	Make two applications 28 days apart in the spring and two applications 28 days apart in the fall.
Zoysia patch (<i>Rhizoctonia solani</i> and/ or <i>Gaeumannomyces</i> <i>incrustana</i>)	0.38-0.77	28	Make one or two applications in late fall before snow cover or when conditions are favorable for disease development. Do not apply on top of snow.

*Do not apply more than two sequential applications of ArmorTech ZOXY 2F for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of ArmorTech ZOXY 2F.

ArmorTech Zoxy 2F Rate Conversion Chart for Turf

Fluid Ounces Product per 1000 sq. ft.	Ounces A.I. per 1000 sq. ft.	Fluid Ounces Product per Acre	Pints of Product per Acre
0.4	0.104	17.4	1.1
0.5	0.130	21.8	1.4
0.6	0.156	26.1	1.6
0.7	0.182	30.5	1.9
0.77	0.200	33.5	2.1
1.35	0.35	58.8	3.7

Amount of ArmorTech Zoxy 2F to Mix 100 Gallons for Turf Applications

ArmorTech Zoxy 2F Use Rate (fl. oz.)	Spray Volume (gallons/1000 square feet)		
	2.0 gals (fl. oz.)	3.0 gals. (fl. oz.)	4.0 gals. (fl. oz.)
0.4	20	13	10
0.5	25	17	13
0.6	30	20	15
0.7	35	23	18
0.77	38.5	25.7	19.3
1.35	67.5	45	33.75

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Turf Alliance, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Turf Alliance, LLC and Seller harmless for any claims relating to such factors.

United Turf Alliance, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Turf Alliance, LLC, and Buyer and User assume the risk of any such use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED TURF ALLIANCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither United Turf Alliance, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED TURF ALLIANCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED TURF ALLIANCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Turf Alliance, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of United Turf Alliance, LLC.

EPA 20140206